

DI

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
28 October 2004 (28.10.2004)

PCT

(10) International Publication Number
WO 2004/091619 A1

(51) International Patent Classification⁷: A61K 31/4709,
C07D 471/04

(21) International Application Number:
PCT/US2004/011031

(22) International Filing Date: 9 April 2004 (09.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
308/MAS/2003 9 April 2003 (09.04.2003) IN

(71) Applicants (for all designated States except US): DR.
REDDY'S LABORATORIES LIMITED [IN/IN];
7-1-27 Ameerpet, Hyderabad, 500 016 Andhra Pradesh
(IN). DR. REDDY'S LABORATORIES, INC. [US/US];
One Park Way, Upper Saddle River, NJ 07458 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): REDDY, Manne,
Satyanarayana [IN/IN]; H.No. 8-3-167/D16, Kalyan Na-
gar-Near AG Colony, Erragadda, Hyderabad-500 038 (IN).
ESWARAJAH, Sajja [IN/IN]; Lig- 110, Dharmareddy
colony-K.P.H.B. Colony, Hyderabad- 500 072 (IN).
RAJU, Vetukuri, Venkata, Naga, Kali, Vara, Prasada

[IN/IN]; Flat No.:404, Shiva durga Residency, H.L.G:
216,217,VI Phase,, K.P.H.B. Colony-Hyderabad 500 072
(IN). KUMAR, Rapolu, Rajesh [IN/IN]; Post:Cherla-
pally, Mandal & Dist., Nalgonda 500 001 (IN). SRINI-
VASREDDY, Ningam [IN/IN]; S/O Ningam Ananthreddy,
Post:Pastapur, Mandal:Zahirabad, Dist:Medak-Pin-502
318 (IN). RAVINDRA, Vedantham [IN/IN]; S/O Vedan-
tham Bhaskara Venkata, D.No.:28-1-80, Narasimha
Charyulu, Kankaragunta-Guntur 522004 (IN).


(74) Agents: CORD, Janet, I. et al.; Ladas & Parry, 26 West
61st Street, New York, NY 10023 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),

[Continued on next page]

(54) Title: A CRYSTALLINE FORM III OF ANHYDROUS MOXIFLOXACIN HYDROCHLORIDE AND A PROCESS FOR
PREPARATION THEREOF



(57) Abstract: A new crystalline Form III of moxifloxacin monohydrochloride and processes for making the crystalline form as well as compositions, pharmaceutical compositions, and methods utilizing the crystalline form are described.

WO 2004/091619 A1

WO 2004/091619 A1



Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

A crystalline Form III of anhydrous moxifloxacin hydrochloride and a process for
preparation thereof

CROSS-REFERENCE TO RELATED APPLICATION

This application claims priority of Indian Patent Application No. 308/MAS/2003, filed April 9, 2003, of which entire content is incorporated herein by reference.

BACKGROUND OF THE INVENTION

Moxifloxacin monohydrochloride is a synthetic broad-spectrum antibacterial agent. The active moiety, moxifloxacin has been shown to be clinically active against most strains of microorganisms such as aerobic gram-positive microorganisms including staphylococcus aureus (methicillin-susceptible strains only), streptococcus pneumoniae (penicillin-susceptible strains), and streptococcus pyogenes; aerobic gram-negative microorganisms including haemophilus influenzae, haemophilus parainfluenzae, klebsiella pneumoniae, and moraxella catarrhalis; and other microorganisms like chlamydia pneumoniae and mycoplasma pneumoniae.

SUMMARY OF INVENTION

In accordance with one aspect, the invention provides a novel crystalline Form III of anhydrous moxifloxacin monohydrochloride. The crystalline Form III of moxifloxacin monohydrochloride may be characterized by an X-ray diffraction pattern, expressed in terms of 2θ angles and obtained with a diffractometer equipped with a copper K X-radiation source, wherein the X-ray powder diffraction pattern includes five or more peaks selected from the group consisting of peaks with 2θ angles of 5.6 ± 0.09 , 7.1 ± 0.09 , 8.4 ± 0.09 , 8.8 ± 0.09 , 10.0 ± 0.09 , 10.4 ± 0.09 , 10.4 ± 0.09 , 11.4 ± 0.09 , 12.2 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 14.4 ± 0.09 , 14.7 ± 0.09 , 16.6 ± 0.09 , 16.9 ± 0.09 , 17.2 ± 0.09 , 17.7 ± 0.09 , 18.5 ± 0.09 , 19.1 ± 0.09 , 19.2 ± 0.09 , 19.8 ± 0.09 , 20.1 ± 0.09 , 20.3 ± 0.09 , 21.1 ± 0.09 , 21.5 ± 0.09 , 22.1 ± 0.09 , 22.6 ± 0.09 , 22.9 ± 0.09 , 23.5 ± 0.09 , 24.0 ± 0.09 , 24.6 ± 0.09 , 24.9 ± 0.09 , 25.8 ± 0.09 , 26.2 ± 0.09 , 26.6 ± 0.09 , 26.9 ± 0.09 , 27.2 ± 0.09 , 28.7 ± 0.09 , 29.1 ± 0.09 , 29.7 ± 0.09 , 30.1 ± 0.09 , 31.4 ± 0.09 , 32.1 ± 0.09 , 37.3 ± 0.09 , 39.0 ± 0.09 , 40.8 ± 0.09 , 41.5 ± 0.09 , 42.2 ± 0.09 , and 43.1 ± 0.09 degrees. The crystalline Form III of moxifloxacin monohydrochloride may also be characterized

by an X-ray diffraction pattern, expressed in terms of 2θ angles and obtained with a diffractometer equipped with a copper K X-radiation source, wherein the X-ray powder diffraction pattern includes two or more peaks selected from the group consisting of peaks with 2θ angles of 7.1 ± 0.09 , 8.8 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 16.6 ± 0.09 , 17.7 ± 0.09 , and 22.1 ± 0.09 . The crystalline form III of anhydrous moxifloxacin monohydrochloride may also be characterized by other analytical methods. Various other embodiments and variants are also provided.

In another aspect, the invention provides a composition that includes moxifloxacin in a solid form, wherein at least 80 % by weight of the solid moxifloxacin monohydrochloride is the crystalline form III of anhydrous moxifloxacin monohydrochloride. The crystalline form III of anhydrous moxifloxacin monohydrochloride in the composition of this aspect of the invention may be characterized by the XRD patterns as described.

The invention also relates to a process for preparing the crystalline form III of moxifloxacin monohydrochloride and to a pharmaceutical composition that includes the crystalline form III of moxifloxacin monohydrochloride and one or more pharmaceutically acceptable carriers or diluents. The pharmaceutical composition may also include one or more additional active ingredients. Preferably, the pharmaceutical composition is in a solid dosage form for oral administration, such as a tablet.

The invention also relates to a method of preventing or treating allergic syndromes, by administering to a patient in need of such treatment an effective amount of crystalline form III of anhydrous moxifloxacin monohydrochloride.

DESCRIPTION OF THE ACCOMPANYING DRAWINGS

Figure 1 is a sample of X-ray powder diffractogram of the crystalline form III of moxifloxacin monohydrochloride.

Figure 2 is a sample of ^{13}C solid state NMR spectrum of the crystalline form III of moxifloxacin monohydrochloride.

Figure 3 is a sample of an infrared spectrum of the crystalline form III of moxifloxacin monohydrochloride.

Figure 4 is a sample of a thermo gravimetric analysis thermogram of the crystalline form III of moxifloxacin monohydrochloride.

Figure 5 is a sample of a differential scanning calorimetry thermogram of the crystalline form III of moxifloxacin monohydrochloride.

DETAILED DESCRIPTION OF THE INVENTION

Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art, to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described.

Unless stated to the contrary, any use of the words such as "including," "containing," "comprising," "having" and the like, means "including without limitation" and shall not be construed to limit any general statement that it follows to the specific or similar items or matters immediately following it. Embodiments of the invention are not mutually exclusive, but may be implemented in various combinations. The described embodiments of the invention and the disclosed examples are given for the purpose of illustration rather than limitation of the invention as set forth the appended claims.

For purposes of the present invention, the following terms are defined below.

A "compound" is a chemical substance that includes molecules of the same chemical structure.

"Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally non-toxic and is not biologically undesirable and includes that which is acceptable for veterinary use and/or human pharmaceutical use.

The term "composition" includes, but is not limited to, a powder, a suspension, an emulsion and/or mixtures thereof. The term composition is intended to encompass a product containing the specified ingredients in the specified amounts, as well as any product, which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. A "composition" may contain a single compound or a mixture of compounds.

The term "pharmaceutical composition" is intended to encompass a product comprising the active ingredient(s), pharmaceutically acceptable excipients that make up the carrier, as well as any product which results, directly or indirectly, from combination,

complexation or aggregation of any two or more of the ingredients, or from dissociation of one or more of the ingredients, or from other types of reactions or interactions of one or more of the ingredients. Accordingly, the pharmaceutical compositions of the present invention encompass any composition made by admixing the active ingredient, additional active ingredient(s), and pharmaceutically acceptable excipients.

The term "excipient" means a component of a pharmaceutical product that is not the active ingredient, such as filler, diluent, carrier, and so on. The excipients that are useful in preparing a pharmaceutical composition are preferably generally safe, non-toxic and neither biologically nor otherwise undesirable, and are acceptable for veterinary use as well as human pharmaceutical use. "A pharmaceutically acceptable excipient" as used in the specification and claims includes both one and more than one such excipient.


The term "isolating" is used to indicate separation of the compound being isolated regardless of the purity of the isolated compound from any unwanted substance which presents with the compound as a mixture. Thus, degree of the purity of the isolated or separated compound does not affect the status of "isolating".

The term "lower alkyl alcohol" as used in the claims defines alcohols having from 1 to 8 carbon atoms and capable of dissolving moxifloxacin monohydrochloride or moxifloxacin at their reflux temperatures with solubility of at least 0.05 g/ml (drug-to-solvent).

The term "substantially free of" in reference to a composition, as used herein, means that the substance cannot be detected in the composition by methods known to those skilled in the art at the time of the filing of this application.

The term "crystalline Form III of anhydrous moxifloxacin monohydrochloride" is used to refer to a new polymorphic form of anhydrous moxifloxacin monohydrochloride obtained by the inventors. In the Indian Patent Application No. 308/MAS/2003, the benefit of priority of which is sought for the present patent application, the substance defined herein as crystalline Form III of anhydrous moxifloxacin monohydrochloride is referred to as "crystalline Form II." The invention contemplates the actual substance of the crystalline Form III of anhydrous moxifloxacin monohydrochloride regardless of its particle size, method of preparation and/or methods of analytical characterization.

Moxifloxacin monohydrochloride is 1-cyclopropyl-7-([S,S]-2,8-diazabicyclo-[4.3.0]non-8-yl)-6-fluoro-1,4-dihydro-8-methoxy-4-oxo-3-quinoline carboxylic acid mono hydrochloride, which is a hydrochloric acid salt of 1-cyclopropyl-7-([S,S]-2,8-diazabicyclo-[4.3.0]non-8-yl)-6-fluoro-1,4-dihydro-8-methoxy-4-oxo-3-quinoline carboxylic acid in 1:1 molar ratio, and has the structure as follows:



As a molecule, moxifloxacin monohydrochloride is described in U.S. 5,607,943, of which entire content is incorporated by reference herein. However, it is known that polymorphic forms of the same drug may have substantial differences in certain pharmaceutically important properties such as dissolution characteristics and bioavailability as well as stability of the drug. Furthermore, difference crystalline form may have different particle size, hardness and glass transition temperature. Thus, one crystalline form may provide significant advantages over other crystalline forms of the same drug in solid dosage form manufacture process such as accurate measurement of the active ingredients, easier filtration, or improved stability during granulation or storage. Furthermore, a particular process suitable for one crystalline form may also provide drug manufacturers several advantages such as economically or environmentally suitable solvents or process, or higher purity or yield of the desired product.

U.S. Patent No. 5,849,752 ("the '752 patent"), incorporated by reference, discloses certain specific crystalline forms of anhydrous moxifloxacin monohydrochloride and monohydrated moxifloxacin monohydrochloride. For convenience, the anhydrous crystalline form disclosed in the '752 patent is designated as

"Form I," and the hydrated form as "Form II." The '752 patent discloses X-ray diffraction patterns of the form I and II as shown in the following table 1:

Table 1

X-ray diffraction patterns (2 θ)	
Anhydrous form I	Hydrated form II
5.8	5.8
8.6	8.5
10.3	10.1
11.6	11.6
13.6	13.4
14.5	14.5
15.0	14.8
15.8	15.6
17.3	17.0
17.5	17.2
18.3	17.4
18.9	17.5
19.3	17.9
19.6	18.6
20.6	19.1
21.5	19.6
22.5	20.4
22.8	21.1
23.0	21.8
23.8	22.7
24.2	23.0
24.7	23.6
25.0	24.1
26.3	24.5
27.0	26.5
27.4	26.7

27.8	27.0
28.2	27.3
29.4	27.5
29.7	27.8
30.0	28.5
30.3	28.9
31.3	29.2
31.8	29.7
34.5	31.4
35.3	31.9
37.1	32.3
	32.6
	34.2
	35.1
	35.5
	36.8
	37.5

Other spectra such ^{13}C solid state NMR, IR, DSC, thermogravimetry and Raman of the form I and II are also disclosed in the '752 patent.

According to one aspect, the present invention provides a new crystalline Form III of anhydrous moxifloxacin monohydrochloride, which is different from the Form I and Form II of the '752 patent. The crystalline Form III of anhydrous moxifloxacin monohydrochloride may be prepared by a process including refluxing azeotropically a mixture of moxifloxacin monohydrochloride and a solvent selected from the group consisting of lower branched or chained acid esters, aliphatic ketones and aliphatic hydrocarbon solvents to form a mixture; cooling the refluxed mixture until solids separate; and isolating said solids thereby obtaining said crystalline form III of moxifloxacin monohydrochloride. Non-limiting examples of the suitable solvents include tertiary butyl acetate, cyclohexane, and toluene.

Alternatively, the crystalline Form III of anhydrous moxifloxacin monohydrochloride may be prepared by a process including dissolving moxifloxacin hydrochloride in a lower alkyl alcohol to obtain a solution; adding to the solution an anti solvent, in which moxifloxacin hydrochloride is poorly soluble but which is miscible with said lower alkyl alcohol; cooling the resulted mixture after adding the anti solvent until solids separate; and isolating said solids thus obtaining the crystalline form III of moxifloxacin monohydrochloride. Preferably, the suitable lower alkyl alcohols with moxifloxacin or moxifloxacin monohydrochloride solubility greater than 0.075 g/ml are used; more preferred lower alkyl alcohols have solubility greater than 0.1 g/ml on drug to solvent basis. The suitable lower alkyl alcohols include, for example, methanol, ethanol, t-butyl alcohol, isopropyl alcohol, other commonly used alkyl alcohol solvents, and mixtures thereof. A suitable non-limiting example of the anti solvent is acetonitrile.

The cooling step of the both processes may be accompanied by stirring the mixtures. The isolation of the solids can be easily done by conventional methods such as filtration, and the isolated compound may be dried at an elevated temperature, which is preferably at about 30-100 °C, more preferably at about 60-90 °C.

In one particular embodiment of the process aspect of the invention, the preparation of the crystalline Form III of anhydrous moxifloxacin hydrochloride includes:

- i) refluxing azeotropically the starting moxifloxacin hydrochloride in lower branched or chained acid esters such as tertiary butyl acetate or an aliphatic hydrocarbon solvent such as cyclohexane or aromatic hydrocarbons such as toluene;
- ii) cooling the reaction mixture of step (i) accompanied by stirring of the mixture till the solid mass crystallizes;
- iii) isolating the solid obtained in step (ii) by conventional methods;

- iv) drying the isolated compound of step (iii) with or without vacuum at 30-100°C, preferably 60-90°C to afford the crystalline Form III of anhydrous moxifloxacin hydrochloride.

In another particular embodiment of the process aspect of the invention, the preparation of the crystalline Form III of anhydrous moxifloxacin hydrochloride includes:

- i) dissolution of the starting moxifloxacin hydrochloride in C₁-C₆ alcohols, such as methanol, at 25-70 °C, preferably at 60-65 °C;
- ii) adding an anti solvents, such as acetonitrile, in which the product is poorly soluble;
- iii) cooling the solution mixture of step (ii) accompanied by stirring of the mixture till the solid mass crystallizes;
- iv) isolating the solid obtained in step (iii) by conventional methods;
- v) drying the isolated compound of step (iv) with or without vacuum at 30-100°C, preferably 60-90°C to afford the crystalline Form III of anhydrous moxifloxacin hydrochloride.

The crystalline form III of moxifloxacin monohydrochloride produced by the inventors was characterized by an X-ray powder diffraction pattern. An example of one X-ray diffraction analysis is shown in Figure 1, and the characteristic 2 theta values (in degrees) in the X-ray diffractograms are shown in Table 2:

Table 2

2 theta (°)
5.6
7.1
8.4
8.8
10.0
10.4

11.4
12.2
13.1
13.9
14.4
14.7
16.6
16.9
17.2
17.7
18.5
19.1
19.2
19.8
20.1
20.3
21.1
21.5
22.1
22.6
22.9
23.5
24.0
24.6
24.9
25.8
26.2
26.6
26.9
27.2
28.7
29.1
29.7
30.1
31.4
32.1
37.3
39.0
40.8
41.5

42.2
43.1

The X-ray diffractogram was measured on a Bruker Axis, D8 Advance Powder X-ray Diffractometer with Cu K alpha-1 radiation source.

It should be kept in mind that slight variations in the observed 2 theta angles values are expected based on the specific diffractometer employed, the analyst and the sample preparation technique. More variation is expected for the relative peak intensities, which is largely affected by the particle size of the sample. Thus, identification of the exact crystalline form of a compound should be based primarily on observed 2 theta angles with lesser importance attributed to relative peak intensities. The 2 theta diffraction angles and corresponding d-spacing values account for positions of various peaks in the X-ray powder diffraction pattern. D-spacing values are calculated with observed 2 theta angles and copper K(α 1) wavelength using the Bragg equation well known to those of skill in the art. Table 3 shows the results of another X-ray diffraction analysis of a sample of crystalline Form III of anhydrous moxifloxacin monohydrochloride, demonstrating the variability in the observed 2 theta angles values:

Table 3

2 theta (°)
5.7
7.1
8.5
8.8
10.0
10.2
10.5
11.4
12.2
13.1
14.0
14.4
14.7
15.1
15.5
16.5
17.2
17.7
18.5
19.2
19.7
20.3
21.6
22.2
23.0
23.6
24.0
24.6
25.0
25.7
26.4
27.2
27.8
28.3
28.9
29.9
32.2
34.9
35.9
36.6
37.3
39.0
41.2
41.8
44.6

Thus, some margin of error may be present in each of the 2 theta angle assignments reported herein. The assigned margin of error in the 2 theta angles for the crystalline form of moxifloxacin monohydrochloride is approximately ± 0.09 for each of the peak assignments. In view of the assigned margin of error, in a preferred variant, the crystalline form III of anhydrous moxifloxacin monohydrochloride may be characterized by an X-ray diffraction pattern, expressed in terms of 2 theta angles, that includes four or more peaks selected from the group consisting of 5.6 ± 0.09 , 7.1 ± 0.09 , 8.4 ± 0.09 , 8.8 ± 0.09 , 10.0 ± 0.09 , 10.4 ± 0.09 , 10.4 ± 0.09 , 11.4 ± 0.09 , 12.2 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 14.4 ± 0.09 , 14.7 ± 0.09 , 16.6 ± 0.09 , 16.9 ± 0.09 , 17.2 ± 0.09 , 17.7 ± 0.09 , 18.5 ± 0.09 , 19.1 ± 0.09 , 19.2 ± 0.09 , 19.8 ± 0.09 , 20.1 ± 0.09 , 20.3 ± 0.09 , 21.1 ± 0.09 , 21.5 ± 0.09 , 22.1 ± 0.09 , 22.6 ± 0.09 , 22.9 ± 0.09 , 23.5 ± 0.09 , 24.0 ± 0.09 , 24.6 ± 0.09 , 24.9 ± 0.09 , 25.8 ± 0.09 , 26.2 ± 0.09 , 26.6 ± 0.09 , 26.9 ± 0.09 , 27.2 ± 0.09 , 28.7 ± 0.09 , 29.1 ± 0.09 , 29.7 ± 0.09 , 30.1 ± 0.09 , 31.4 ± 0.09 , 32.1 ± 0.09 , 37.3 ± 0.09 , 39.0 ± 0.09 , 40.8 ± 0.09 , 41.5 ± 0.09 , 42.2 ± 0.09 , and 43.1 ± 0.09 degrees.

Comparing the XRD data in Tables 1, 2, and 3, it is apparent that certain peaks provide the best way of characterizing the crystalline Form III of anhydrous moxifloxacin monohydrochloride and of differentiating it from the Forms I and II. Very few of such peaks are needed to allow for such characterization and differentiation, including presence of the crystalline Form III of anhydrous moxifloxacin monohydrochloride in mixtures with other forms of moxifloxacin. Thus, the crystalline Form III of anhydrous moxifloxacin monohydrochloride may also be characterized by an X-ray powder diffraction pattern includes two or more peaks selected from the group consisting of peaks with 2 theta angles of 7.1 ± 0.09 , 8.8 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 16.6 ± 0.09 , 17.7 ± 0.09 , and 22.1 ± 0.09 .

Since some margin of error is possible in the assignment of 2 theta angles and d-spacings, the preferred method of comparing X-ray powder diffraction patterns in order to identify a particular crystalline form is to overlay the X-ray powder diffraction pattern of the unknown form over the X-ray powder diffraction pattern of a known form. For example, one skilled in the art can overlay an X-ray powder diffraction pattern of an unidentified crystalline form of moxifloxacin monohydrochloride over Fig. 1 and readily determine whether the X-ray diffraction pattern of the unidentified form is substantially

the same as the X-ray powder diffraction pattern of the crystalline form of this invention. If the X-ray powder diffraction pattern is substantially the same as FIG. 1, the previously unknown crystalline form of moxifloxacin monohydrochloride can be readily and accurately identified as the crystalline Form III of this invention.

The crystalline Form III of moxifloxacin monohydrochloride is anhydrous. A sample of the crystalline Form III prepared by the inventors had moisture content less than 0.2 % by KF method, which confirmed the anhydrous nature of the compound. While the invention is not limited to any specific theory, it should be understood however that the crystalline form III of moxifloxacin monohydrochloride may contain residual, unbound moisture without losing its anhydrous character and/or its crystalline form III characteristics. It is believed that residual moisture may be present in the form of water molecules in the channel of the crystals, rather than being bound inside the crystal lattice as in hydrated forms. When the anhydrous crystalline form is wet, the entire crystalline lattice may expand due to the space occupied by the water molecules. Then the X-ray powder diffraction pattern of the wet crystalline form may also expand. In such case, the X-ray powder diffraction patterns of two different moisture content crystalline forms may not be perfectly overlapped. Nevertheless, one of skill in the art should be able to determine whether they are same crystalline forms or not, by looking at the overall shape of the X-ray powder diffraction pattern optionally with help of other spectroscopy data such as Infrared spectroscopy (IR).

The crystalline form III of moxifloxacin monohydrochloride prepared by the inventors was also characterized by ^{13}C solid state NMR and IR as shown respectively in Figs. 2 and 3. The NMR spectrum includes a characteristic peak at about 107 ppm. Table 4 shows a comparison between the ^{13}C solid state NMR spectra of Forms I, II, and III:

Table 4:

Form I	Form II	Form III
	7.7	
	8.3	
8.5		
	9.0	
	10.8	
		11.612
	12.1	
12.3		

14.1		
		14.792
18.2	18.2	
	19.8	
20.2		
		20.465
22.8	22.9	
		25.013
	34.9	
35.2		
		37.390
39.7		
	40.2	
		42.312
46.5		
	47.0	
		48.836
49.5	49.5	
	50.1	
52.3		
		52.431
	52.6	
		54.443
55.9	55.9	
	56.8	
		57.792
59.2		
	59.4	
		61.240
62.6		
	64.1	
		65.261
65.8		
	66.8	
		29.240
	105.0	
105.4		
	107.1	107.100
108.1		
		109.157
		110.303
	116.3	
116.9		
117.5	117.4	
		117.687
		120.043
134.7		
	135.2	
136.0	136.1	
137.3	137.4	137.395
		139.383
140.1		

	140.8	
		142.188
142.6		
	143.5	
		145.897
	149.3	
150.1		
	150.9	
152.6		
		153.516
165.3		
166.0		166.597
	168.1	167.918
		171.145
175.5	175.5	
		177.275

In the IR spectrum, the peak locations of several distinctive peaks may help one of skill in the art to identify the crystalline form of the present invention. These peaks include absorption bands at about 1159 cm^{-1} , 1459 cm^{-1} , about 1515 cm^{-1} , about 1623 cm^{-1} , and 2706 cm^{-1} . The ^{13}C solid state NMR spectrum was measured with a Bruker MSL 300, and the IR spectrum was measured by KBr-transmission method with Perkin Elmer IR spectroscopy.

The Differential scanning calorimetry (DSC) thermogram of crystalline form of Moxifloxacin monohydrochloride obtained by the inventors is shown in Figure 5. It exhibits a significant endo-exo pattern with identified peaks around $246\text{ }^{\circ}\text{C}$. The DSC spectrum was measured on a Perkin Elmer Pyris 6 DSC. It is known to one of skill in the art that the endothermic peak location may be affected by the heating rate in the DSC. Thus, slight variation of the peak may be acceptable.

The invention also relates to a composition containing solid moxifloxacin monohydrochloride of which at least 80%, by total weight of the solid moxifloxacin monohydrochloride in the composition, is the crystalline form III. In the more preferred form of this composition, the solid moxifloxacin monohydrochloride is suitable for use as active ingredient in formulating pharmaceutical products. In an embodiment of the invention, the composition may comprise at least 90% of the crystalline form III of moxifloxacin monohydrochloride with respect to total weight of the solid moxifloxacin monohydrochloride in the composition. In another embodiment of the invention, the composition may comprise at least 95% of the crystalline form III of moxifloxacin

monohydrochloride with respect to total weight of the solid moxifloxacin monohydrochloride in the composition. In yet another embodiment of the invention, the composition is substantially free of the form I and II of moxifloxacin monohydrochloride.

X-ray diffraction provides a convenient and practical means for quantitative determination of the relative amounts of crystalline and/or amorphous forms in a solid mixture. X-ray diffraction is adaptable to quantitative applications because the intensities of the diffraction peaks, particularly long range peaks of a given compound in a mixture are proportional to the fraction of the corresponding powder in the mixture. The percent composition of crystalline moxifloxacin monohydrochloride in an unknown composition can be determined. Preferably, the measurements are made on solid powder moxifloxacin monohydrochloride. The X-ray powder diffraction patterns of an unknown composition can be compared to known quantitative standards containing the pure crystalline form III of moxifloxacin monohydrochloride to identify the percent ratio of a particular crystalline form. This is done by comparing the relative intensities of the peaks from the diffraction pattern of the unknown solid powder composition with a calibration curve derived from the X-ray diffraction patterns of pure known samples. The curve can be calibrated based on the X-ray powder diffraction pattern for the strongest peak or any distinctive peak from a pure sample of the crystalline form III of moxifloxacin monohydrochloride. The calibration curve may be created in a manner known to those of skill in the art. For example, five or more artificial mixtures of crystalline forms of moxifloxacin monohydrochloride, at different amounts, may be prepared. In a non-limiting example, such mixtures may contain, 2%, 5%, 7%, 8%, and 10% of the crystalline III moxifloxacin monohydrochloride. Then, X-ray diffraction patterns are obtained for each artificial mixture using standard X-ray diffraction techniques. Slight variations in peak positions, if any, may be accounted for by adjusting the location of the peak to be measured. The intensities of the selected characteristic peak(s) for each of the artificial mixtures are then plotted against the known weight percentages of the crystalline form. The resulting plot is a calibration curve that allows determination of the amount of the crystalline form III of moxifloxacin monohydrochloride in an unknown sample. For the unknown mixture of the crystalline and amorphous forms of moxifloxacin monohydrochloride, the intensities of the selected characteristic peak(s) in

the mixture, relative to an intensity of this peak in a calibration mixture, may be used to determine the percentage of the given crystalline form in the composition, with the remainder determined to be the amorphous material.

Similar quantitative analysis may be done using IR spectroscopy, particularly with attenuating total reflectance (ATR) technology.

Pharmaceutical compositions comprising crystalline form III of moxifloxacin monohydrochloride can be formulated with one or more pharmaceutically acceptable carriers, also known as excipients, which ordinarily lack pharmaceutical activity, but have various useful properties which may, for example, enhance the stability, sterility, bioavailability, and ease of formulation of a pharmaceutical composition. These carriers are pharmaceutically acceptable, meaning that they are not harmful to humans or animals when taken appropriately and are compatible with the other ingredients in a given formulation. The carriers may be solid, semi-solid, or liquid, and may be formulated with the compound in bulk. The resulting mixture may be manufactured in the form of a unit-dose formulation (i.e., a physically discrete unit containing a specific amount of active ingredient) such as a tablet or capsule.

Generally, the pharmaceutical compositions of the invention may be prepared by uniformly admixing the active ingredient with liquid or solid carriers and then shaping the product into the desired form. The pharmaceutical compositions may be in the form of suspensions, solutions, elixirs, aerosols, or solid dosage forms. Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are employed.

A preferred oral solid preparation is a tablet. A tablet may be prepared by direct compression, wet granulation, or molding, of the active ingredient(s) with a carrier and other excipients in a manner known to those skilled in the art. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active agent or dispersing agent. Molded tablets may be made on a suitable machine. A mixture of the powdered compound moistened with an inert liquid diluent is suitable in the case of oral solid dosage forms (e.g., powders, capsules, and tablets). If desired, tablets may be coated by standard techniques. The compounds of this invention

may be formulated into typical disintegrating tablets, or into controlled or extended release dosage forms.

The pharmaceutical compositions of the invention are contemplated in various formulations suitable for various modes of administration, including but not limited to inhalation, oral, rectal, parenteral (including subcutaneous, intradermal, intramuscular, intravenous), implantable, intravaginal and transdermal administration. The most suitable route of administration in any given case depends on the duration of the subject's condition, the length of treatment desired, the nature and severity of the condition being treated, and the particular formulation that is being used. The formulations may be in bulk or in unit dosage form.

The amount of active ingredient included in a unit dosage form depends on the type of formulation that is formulated. A pharmaceutical composition of the invention will generally comprise about 0.1% by weight to about 99% by weight of active ingredient, preferably about 1% by weight to 50% by weight for oral administration and about 0.2% by weight to about 20% by weight for parenteral administration.

Formulations suitable for oral administration include capsules (hard and soft), cachets, lozenges, syrups, suppositories, and tablets, each containing a pre-determined amount of the active compound; as a powder or granules; as a solution or a suspension in an aqueous or non-aqueous liquid; or as an oil-in-water or water-in-oil emulsion. Such formulations may be prepared by any suitable method of pharmacy that includes the step of bringing into association the active compound and a suitable carrier or carriers. For liquid oral formulations, a preferable amount is from about 2% by weight to about 20% by weight. Suitable carriers include but are not limited to fillers, binders, lubricants, inert diluents, surface active/dispersing agents, flavorants, antioxidants, bulking and granulating agents, adsorbants, preservatives, emulsifiers, suspending and wetting agents, glidants, disintegrants, buffers and pH-adjusting agents, and colorants. Examples of carriers include celluloses, modified celluloses, cyclodextrins, starches, oils, polyols, sugar alcohols and sugars, and others. For liquid formulations sugar, sugar alcohols, ethanol, water, glycerol, and polyalkylene glycols are particularly suitable, and may also be used in solid formulations. Cyclodextrins may be particularly useful for increasing bioavailability. Formulations for oral administration may optionally include enteric

coatings known in the art to prevent degradation of the formulation in the stomach and provide release of the drug in the small intestine.

Formulations suitable for buccal or sub-lingual administration include lozenges comprising the active compound in a flavored base, usually sucrose and acacia or tragacanth, although other agents are also suitable, and pastilles comprising the compound in an inert base such as gelatin and glycerin or sucrose and acacia.

Formulations suitable for rectal administration are preferably presented as unit dose suppositories. These may be prepared by admixing the active compound with one or more conventional solid carriers, e.g., cocoa butter, and then shaping the resulting mixture.

In another aspect, the invention also provides methods of treating infections caused by susceptible strains of streptococcus pneumoniae, haemophilus influenzae, moraxella catarrhalis, haemophilus parainfluenzae, klebsiella pneumoniae, staphylococcus aureus, mycoplasma pneumoniae, Chlamydia pneumoniae and streptococcus pyogenes, which includes administering a mammal in need thereof an effective amount of the crystalline form III of moxifloxacin monohydrochloride.

The effective amount (i.e., dosage) of active compound for treatment will vary depending on the route of administration, the condition being treated, its severity, and duration, and the state and age of the subject. A skilled physician will monitor the progress of the subject and will adjust the dosage accordingly, depending on whether the goal is to eliminate, alleviate, or prevent a given condition. Generally, the dosage should be considered in proportion to the subject's weight. The daily dose of particular formulations of active compound may be divided among one or several unit dose administrations. For example therapeutic administration about fifteen to thirty minutes before main meals is preferable (i.e. three times daily), although administration of the active compounds may be carried out prophylactically, and may be maintained for prolonged periods of time. One skilled in the art will take such factors into account when determining dosage. Unit dosage of active ingredient may range preferably from about 1mg to about 800 mg, more preferably from about 100 mg to about 600 mg, even more preferably from about 300 mg to about 500 mg.

The invention is further described by reference to the following examples which set forth in detail the preparation of compounds and compositions of the present invention, as well as their utility. It will be apparent to those skilled in the art, that many modifications, both to materials, and methods, may be practiced without departing from the purpose and interest of this invention. The examples that follow are not intended to limit the scope of the invention as described hereinabove or as claimed below.

Reference Example. Preparation of Moxifloxacin

1-cyclopropyl -6, 7-difluoro-1, 4-dihydro-4-oxo-8-methoxy quinolone-3-carboxylic acid (100 grams), (S,S) Diazabicyclo nonane (60 grams) and 1,8-Diazabicyclo[5.4.0.]undec-7-ene(DBU) (10.gms) were added to N-Methylpyrrolidinone (250 ml) and the reaction mixture was slowly heated to the 60-70°C temperature and stirred till the reaction was substantially completed. 5% aqueous isopropyl alcohol was added to the reaction mass, and pH was adjusted towards basic with caustic lye. Then the reaction mass was filtered, through clarifying filter, washed with 5% aqueous isopropyl alcohol. Combined total filtrate and adjusted pH to 7.0 to 7.2 with aqueous Hcl. and isolated at a temperature of 10-15°C to afford moxifloxacin. Moxifloxacin then treated with Hydrochloric acid in 10% aqueous methanol to yield corresponding hydrochloride salt. (Wet weight: 115 grams)

EXAMPLE 1. Preparation of novel crystalline form III of moxifloxacin hydrochloride.

Moxifloxacin hydrochloride (50 grams) (obtained from reference example) was suspended in tertiary butyl acetate (250 ml) and heated to reflux temperature of 90-100°C. Water was azeotropically removed, accompanied by cooling the reaction mixture to a temperature of 10-15°C under stirring for 30-60 mints to crystallize the solid mass. The crystallized mass was filtered, and washed with tertiary butyl acetate (50 ml) and dried at a temperature of 60-70°C to afford the novel crystalline form III of moxifloxacin hydrochloride. (Weight: 46.8 grams, M.C. by KF is 0.20%)

EXAMPLE 2. Preparation of novel crystalline form III of anhydrous moxifloxacin hydrochloride

Moxifloxacin Hydrochloride (115 grams) (obtained as per reference example) was dissolved in methanol (1000 ml) at reflux temperature accompanied by gently stirring for 30 min. Acetonitrile (1500 ml) was added to the above solution, the resultant

solution was cooled to a temperature of 25-35°C and stirred for 21 hrs. The obtained solid mass was filtered and dried at a temperature of 50-70°C to afford the Novel crystalline form III of anhydrous Moxifloxacin hydrochloride.

(Weight: 49 grams, M.C. by KF is 0.2%)

EXAMPLE 3. Preparation of novel crystalline form III of anhydrous moxifloxacin hydrochloride

Moxifloxacin Hydrochloride (40 grams) (obtained from reference example) was suspended in methyl isobutyl ketone (400 ml) and heated to 110-120°C, while collecting the low boilers and refluxed azeotropically between 115-120°C and then reaction mass is cooled to 25-35°C and product is filtered and dried at 80-90°C under vacuum to afford the novel crystalline form III of anhydrous Moxifloxacin hydrochloride.

(Weight: 35.8gms, M.C. by KF is 0.20%; Purity by HPLC: 99.88%).

EXAMPLE 4. Preparation of novel crystalline form III of anhydrous moxifloxacin hydrochloride

1-cyclopropyl -6, 7-difluoro-1, 4-dihydro-4-oxo-8-methoxy quinolone-3-carboxylic acid (50 Kgs), (S,S) diazabicyclo nonane (1.49 equivalents) and 1,8-diazabicyclo [5.4.0.] undec-7-ene(DBU) (5Kgs) were added to N-methylpyrrolidinone (125L) in SS Reactor and the reaction mixture was slowly heated to the 60-65°C temperature and stirred till the reaction was substantially completed. 500L of 5% aqueous isopropyl alcohol was added to the reaction mass, and pH was adjusted to 5.0-6.0 and the product is isolated at 20-25°C. Wet cake is recrystallised in aqueous methanol at pH 1.5-2.0, and is made slurry in 5% aqueous methanol. Then wet cake was dissolved in aqueous methanol and the reaction mass was filtered through clarifying filter, washed with aqueous methanol. Combined total filtrate pH was adjusted to 1.5-2.0 with Aqueous HCl. Finally, wet cake is taken with methyl isobutylketone (800ml) and heated to reflux while collecting the low boilers and refluxed azeotropically between 115-120°C and then reaction mass is cooled to 25-35°C and product is filtered and dried at 80-90°C under vacuum to afford the novel crystalline form III of anhydrous moxifloxacin hydrochloride. (Weight: 31.3Kgs, M.C. by KF is 0.60%; Purity by HPLC: 99.94%).

Example 5. Soluble granules containing the crystalline form III of moxifloxacin monohydrochloride.

Soluble granules containing crystalline moxifloxacin monohydrochloride may have the following content:

Ingredient	Content (mg)
Crystalline Form III of anhydrous moxifloxacin monohydrochloride	400
Calcium carbonate	800
Citric acid	900
Avicel	40
Mannitol	625
Maltodextrin	15
Aspartame	3
Aroma	20

Example 6. Dispersible tablet containing crystalline moxifloxacin monohydrochloride.

Dispersible tablet containing crystalline moxifloxacin monohydrochloride may have the following content:

Ingredient	Content (mg)
Crystalline Form III of anhydrous moxifloxacin monohydrochloride	400
Calcium carbonate	500
Polyvinylpyrrolidone	17
Avicel	15
Mannitol	400
Maltodextrin	15
Aspartame	3
Aroma	20

EXAMPLE 7. A tablet containing anhydrous monofloxacin hydrochloride

A tablet containing crystalline moxifloxacin monohydrochloride had the following content:

Ingredient	Quantity/tab(mg)
Slug composition	
Drug Premix	
Moxifloxacin HCl (Anhydrous)	436.30
Colloidal Silicon Dioxide	7.05
Talc	4.70
Magnesium Stearate	1.50
EXCIPIENTS	95.03
Microcrystalline Cellulose (Avicel PH 112)	140.85
Lactose Monohydrate (Pharmatose DCL - 21)	26.00
Croscarmellose Sodium	24.00
Talc	2.35
Magnesium Stearate	3.50
Colloidal Silicon Dioxide	1.75
	650.00
Lubrication of milled & sieved slugs	
Croscarmellose Sodium	40.00
Microcrystalline Cellulose (Avicel PH 112)	24.00
Talc	6.00
Magnesium Stearate	3.50
Colloidal Silicon Dioxide	6.50
Tablet Wt.	730
Film Coating (3%w/w)	
Opadry Beige YS-1-17174-A	21.90
Isopropyl Alcohol (70%)	q.s
Methylene Chloride (30%)	q.s
Tablet weight	752.00

Unless stated to the contrary, any use of the words such as "including," "containing," "comprising," "having" and the like, means "including without limitation" and shall not be construed to limit any general statement that it follows to the specific or similar items or matters immediately following it. Except where the context indicates to

the contrary, all exemplary values are intended to be fictitious, unrelated to actual entities and are used for purposes of illustration only. Most of the foregoing alternative embodiments are not mutually exclusive, but may be implemented in various combinations. As these and other variations and combinations of the features discussed above can be utilized without departing from the invention as defined by the claims, the foregoing description of the embodiments should be taken by way of illustration rather than by way of limitation of the invention as defined by the appended claims.

We claim:

1. A compound which is a crystalline Form III of anhydrous moxifloxacin monohydrochloride.

2. The compound of claim 1 having an X-ray diffraction pattern, expressed in terms of 2θ angles and obtained with a diffractometer equipped with a copper K X-radiation source, wherein said X-ray powder diffraction pattern includes five or more peaks selected from the group consisting of peaks with 2θ angles of 5.6 ± 0.09 , 7.1 ± 0.09 , 8.4 ± 0.09 , 8.8 ± 0.09 , 10.0 ± 0.09 , 10.4 ± 0.09 , 10.4 ± 0.09 , 11.4 ± 0.09 , 12.2 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 14.4 ± 0.09 , 14.7 ± 0.09 , 16.6 ± 0.09 , 16.9 ± 0.09 , 17.2 ± 0.09 , 17.7 ± 0.09 , 18.5 ± 0.09 , 19.1 ± 0.09 , 19.2 ± 0.09 , 19.8 ± 0.09 , 20.1 ± 0.09 , 20.3 ± 0.09 , 21.1 ± 0.09 , 21.5 ± 0.09 , 22.1 ± 0.09 , 22.6 ± 0.09 , 22.9 ± 0.09 , 23.5 ± 0.09 , 24.0 ± 0.09 , 24.6 ± 0.09 , 24.9 ± 0.09 , 25.8 ± 0.09 , 26.2 ± 0.09 , 26.6 ± 0.09 , 26.9 ± 0.09 , 27.2 ± 0.09 , 28.7 ± 0.09 , 29.1 ± 0.09 , 29.7 ± 0.09 , 30.1 ± 0.09 , 31.4 ± 0.09 , 32.1 ± 0.09 , 37.3 ± 0.09 , 39.0 ± 0.09 , 40.8 ± 0.09 , 41.5 ± 0.09 , 42.2 ± 0.09 , and 43.1 ± 0.09 degrees.

3. The compound of claim 2, wherein said X-ray diffraction pattern includes peaks with 2θ angles of about 5.6, 7.1, 8.4, 8.8, 10.0, 10.4, 10.4, 11.4, 12.2, 13.1, 13.9, 14.4, 14.7, 16.6, 16.9, 17.2, 17.7, 18.5, 19.1, 19.2, 19.8, 20.1, 20.3, 21.1, 21.5, 22.1, 22.6, 22.9, 23.5, 24.0, 24.6, 24.9, 25.8, 26.2, 26.6, 26.9, 27.2, 28.7, 29.1, 29.7, 30.1, 31.4, 32.1, 37.3, 39.0, 40.8, 41.5, 42.2, and 43.1 degrees.

4. The compound of claim 1 having an X-ray diffraction pattern, expressed in terms of 2θ angles and obtained with a diffractometer equipped with a copper K X-radiation source, wherein said X-ray powder diffraction pattern includes two or more peaks selected from the group consisting of peaks with 2θ angles of 7.1 ± 0.09 , 8.8 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 16.6 ± 0.09 , 17.7 ± 0.09 , and 22.1 ± 0.09 .

5. The compound of claim 1 having substantially the same X-ray diffraction pattern as shown in Figure 1.

6. The compound of claim 1 having a ^{13}C solid state NMR spectrum comprising a peak at about 107 ppm.

7. The compound of claim 1 having substantially the same ^{13}C solid-state NMR spectrum as shown in Figure 2.

8. The compound of claim 1 having an infrared absorption spectrum comprising absorption bands at about 1159 cm^{-1} and 2706 cm^{-1} .

9. The compound of claim 1 having substantially the same infrared spectrum as shown in Figure 3.

10. The compound of claim 1 having a differential scanning calorimetry thermogram, which exhibits an endotherm peak at about $246\text{ }^{\circ}\text{C}$.

11. The compound of claim 1 having substantially the same differential scanning calorimetry thermogram as shown in Figure 5.

12. The compound of claim 1 having substantially the same analytical characterization data as shown in Figures 1, 2, 3, 4, and 5.

13. A composition comprising moxifloxacin monohydrochloride as a solid, wherein at least 80% by weight of said solid moxifloxacin monohydrochloride is a crystalline form III of anhydrous moxifloxacin monohydrochloride having an X-ray diffraction pattern, expressed in terms of 2θ angles and obtained with a diffractometer equipped with a copper K X-radiation source, wherein said X-ray powder diffraction pattern includes five or more peaks selected from the group consisting of peaks with 2θ angles of 5.6 ± 0.09 , 7.1 ± 0.09 , 8.4 ± 0.09 , 8.8 ± 0.09 , 10.0 ± 0.09 , 10.4 ± 0.09 , 10.4 ± 0.09 , 11.4 ± 0.09 , 12.2 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 14.4 ± 0.09 , 14.7 ± 0.09 , 16.6 ± 0.09 , 16.9 ± 0.09 , 17.2 ± 0.09 , 17.7 ± 0.09 , 18.5 ± 0.09 , 19.1 ± 0.09 , 19.2 ± 0.09 , 19.8 ± 0.09 , 20.1 ± 0.09 , 20.3 ± 0.09 , 21.1 ± 0.09 , 21.5 ± 0.09 , 22.1 ± 0.09 , 22.6 ± 0.09 , 22.9 ± 0.09 , 23.5 ± 0.09 , 24.0 ± 0.09 , 24.6 ± 0.09 , 24.9 ± 0.09 , 25.8 ± 0.09 , 26.2 ± 0.09 , 26.6 ± 0.09 , 26.9 ± 0.09 , 27.2 ± 0.09 , 28.7 ± 0.09 , 29.1 ± 0.09 , 29.7 ± 0.09 , 30.1 ± 0.09 , 31.4 ± 0.09 , 32.1 ± 0.09 , 37.3 ± 0.09 , 39.0 ± 0.09 , 40.8 ± 0.09 , 41.5 ± 0.09 , 42.2 ± 0.09 , and 43.1 ± 0.09 degrees.

14. A composition comprising moxifloxacin monohydrochloride as a solid, wherein at least 80% by weight of said solid moxifloxacin monohydrochloride is a crystalline Form III of anhydrous moxifloxacin monohydrochloride having an X-ray diffraction pattern, expressed in terms of 2θ angles and obtained with a diffractometer equipped with a copper K X-radiation source, wherein said X-ray diffraction pattern includes two or more peaks selected from the group consisting of peaks with 2θ

angles of 7.1 ± 0.09 , 8.8 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 16.6 ± 0.09 , 17.7 ± 0.09 , and 22.1 ± 0.09 .

15. The composition of claim 14, wherein at least 90% by weight of said solid moxifloxacin monohydrate is in said crystalline Form III.

16. The composition of claim 14, wherein at least 95% by weight of said solid moxifloxacin monohydrate is in said crystalline Form III.

17. The composition of claim 14, wherein at least 99% by weight of said solid moxifloxacin monohydrate is in said crystalline Form III.

18. A pharmaceutical composition, which comprises a pharmaceutically effective amount of a crystalline Form III of anhydrous moxifloxacin monohydrochloride and one or more pharmaceutically acceptable carriers or diluents.

19. The pharmaceutical composition of claim 18, wherein said crystalline form III of anhydrous moxifloxacin monohydrochloride has an X-ray diffraction pattern expressed in terms of 2 theta angles and obtained with a copper K X-radiation source, wherein said X-ray powder diffraction pattern includes five or more peaks selected from the group consisting of peaks with 2 theta angles of 5.6 ± 0.09 , 7.1 ± 0.09 , 8.4 ± 0.09 , 8.8 ± 0.09 , 10.0 ± 0.09 , 10.4 ± 0.09 , 10.4 ± 0.09 , 11.4 ± 0.09 , 12.2 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 14.4 ± 0.09 , 14.7 ± 0.09 , 16.6 ± 0.09 , 16.9 ± 0.09 , 17.2 ± 0.09 , 17.7 ± 0.09 , 18.5 ± 0.09 , 19.1 ± 0.09 , 19.2 ± 0.09 , 19.8 ± 0.09 , 20.1 ± 0.09 , 20.3 ± 0.09 , 21.1 ± 0.09 , 21.5 ± 0.09 , 22.1 ± 0.09 , 22.6 ± 0.09 , 22.9 ± 0.09 , 23.5 ± 0.09 , 24.0 ± 0.09 , 24.6 ± 0.09 , 24.9 ± 0.09 , 25.8 ± 0.09 , 26.2 ± 0.09 , 26.6 ± 0.09 , 26.9 ± 0.09 , 27.2 ± 0.09 , 28.7 ± 0.09 , 29.1 ± 0.09 , 29.7 ± 0.09 , 30.1 ± 0.09 , 31.4 ± 0.09 , 32.1 ± 0.09 , 37.3 ± 0.09 , 39.0 ± 0.09 , 40.8 ± 0.09 , 41.5 ± 0.09 , 42.2 ± 0.09 , and 43.1 ± 0.09 degrees.

20. The pharmaceutical composition of claim 18, wherein said crystalline Form III of anhydrous moxifloxacin monohydrochloride has an X-ray diffraction pattern expressed in terms of 2 theta angles and obtained with a copper K X-radiation source, wherein said X-ray powder diffraction pattern includes five or more peaks selected from the group consisting of peaks with 2 theta angles of 7.1 ± 0.09 , 8.8 ± 0.09 , 13.1 ± 0.09 , 13.9 ± 0.09 , 16.6 ± 0.09 , 17.7 ± 0.09 , and 22.1 ± 0.09 .

21. The pharmaceutical composition of claim 18, which is a solid dosage form for an oral administration.

22. The pharmaceutical composition of claim 18, wherein said solid dosage form is a tablet.

23. The pharmaceutical composition of claim 18, which is in dosage unit form containing from about 0.5 to about 800 mg of moxifloxacin monohydrochloride.

24. A process for preparation of a crystalline Form III of moxifloxacin monohydrochloride, said process comprising:

a) refluxing azeotropically a starting moxifloxacin monohydrochloride in a solvent selected from the group consisting of lower branched esters, chained acid esters, aliphatic ketones and aliphatic hydrocarbon solvents;

b) cooling the refluxed solvent while stirring the mixture until a solid separates; and

c) isolating said separated solid thereby obtaining said crystalline Form III of anhydrous moxifloxacin monohydrochloride.

25. The process of claim 24, wherein said solvent is selected from the group consisting of tertiary butyl acetate, cyclohexane, toluene, methylisobutylketone, and mixtures thereof.

26. A process for preparation of a crystalline Form III of moxifloxacin monohydrochloride, said process comprising:

a) dissolving moxifloxacin hydrochloride in a lower alkyl alcohol to obtain a solution;

b) adding to the solution an anti solvent, in which moxifloxacin hydrochloride is poorly soluble;

c) cooling the mixed solvents until a solid separates; and

d) isolating said solids thereby obtaining said crystalline Form III of moxifloxacin monohydrochloride.

27. The process of claim 26, wherein said lower alkyl alcohol is selected from the group consisting of methanol, ethanol, t-butyl alcohol, isopropyl alcohol and mixtures thereof.

28. The process of claim 26, wherein said lower alkyl alcohol is methanol.

29. The process of claim 26, wherein said anti solvent is acetonitrile.

30. The moxifloxacin monohydrochloride produced in accordance with the process of claim 24.


31. The moxifloxacin monohydrochloride produced in accordance with the process of claim 26.

32. A method of treating infections caused by susceptible strains of streptococcus pneumoniae, haemophilus influenzae, moraxella catarrhalis, haemophilus parainfluenzae, klebsiella pneumoniae, staphylococcus aureus, mycoplasma pneumoniae, Chlamydia pneumoniae and streptococcus pyogenes, which comprises administering a mammal in need thereof an effective amount of the compound of claim 1.

33. The method of claim 28, wherein said mammal is a human.

1/5

FIG. 1



2/5

FIG. 2

MF 005H03, ¹³C cpmas
at 5KHz






FIG. 3



4/5


FIG. 4

Thermal Analysis Data



5/5

FIG. 5



INTERNATIONAL SEARCH REPORT

International Application No

PCT/US2004/011031

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/4709 C07D471/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, BIOSIS, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 849 752 A (GRUNENBERG ALFONS ET AL) 15 December 1998 (1998-12-15) cited in the application claims 1-10; tables 1-5	1-33
A	WO 99/26940 A (DIEHL HERBERT ; MOHRS KLAUS (DE); BAYER AG (DE); HEILMANN WERNER (DE);) 3 June 1999 (1999-06-03) examples 1,2	1-33

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

12 August 2004

Date of mailing of the international search report

20/08/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Stroeter, T

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2004/011031

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 32 and 33 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/011031

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5849752	A	15-12-1998	DE 19546249 A1	19-06-1997
			AT 221531 T	15-08-2002
			AU 708006 B2	29-07-1999
			AU 7421696 A	19-06-1997
			BG 62258 B1	30-06-1999
			BG 101043 A	30-04-1998
			BR 9605968 A	18-08-1998
			CA 2192418 A1	13-06-1997
			CN 1160052 A , B	24-09-1997
			CZ 9603646 A3	16-07-1997
			DE 59609501 D1	05-09-2002
			DK 780390 T3	11-11-2002
			EE 9600201 A	16-06-1997
			EP 0780390 A1	25-06-1997
			ES 2179910 T3	01-02-2003
			HR 960558 A1	28-02-1998
			HU 9603428 A2	28-08-1997
			ID 22625 A	02-12-1999
			IL 119795 A	27-12-1998
			IN 185805 A1	05-05-2001
			JP 9169757 A	30-06-1997
			MA 24342 A1	01-07-1998
			NL 300109 I1	03-02-2003
			NO 965298 A	13-06-1997
			NZ 299905 A	24-09-1998
			PL 317415 A1	23-06-1997
			PT 780390 T	29-11-2002
			RU 2162468 C2	27-01-2001
			SG 47201 A1	20-03-1998
			SI 780390 T1	31-10-2002
			SK 159196 A3	08-10-1997
			TR 970481 A2	21-06-1997
			TW 411340 B	11-11-2000
			ZA 9610405 A	23-06-1997
WO 9926940	A	03-06-1999	DE 19751948 A1	27-05-1999
			AU 732977 B2	03-05-2001
			AU 1561999 A	15-06-1999
			BG 104467 A	31-08-2001
			BR 9814894 A	03-10-2000
			CA 2311540 A1	03-06-1999
			CN 1418879 A	21-05-2003
			CN 1279683 T	10-01-2001
			EE 200000241 A	15-06-2001
			WO 9926940 A2	03-06-1999
			EP 1034173 A2	13-09-2000
			HR 20000332 A1	30-04-2001
			HU 0004337 A2	28-10-2001
			ID 26840 A	15-02-2001
			JP 2001524477 T	04-12-2001
			NO 20002637 A	23-05-2000
			NZ 504657 A	27-04-2001
			PL 341088 A1	26-03-2001
			RU 2219175 C2	20-12-2003
			SK 7482000 A3	09-10-2000
			TR 200001472 T2	21-06-2002
			TW 513427 B	11-12-2002
			US 2003208069 A1	06-11-2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US2004/011031

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9926940	A	ZA 9810669 A	26-05-1999

